

The following Tables identify the proportion of clients prevention counseled in 1999 who indicated a specific drug use during sexual or needle sharing activities in the previous year.

Tarrant

White

| | M/MS | IDU | | F/MS | |
|--------------|-------|-------|--------|-------|--------|
| | Male | Male | Female | Male | Female |
| Drug use | 57.4% | 89.7% | 88.1% | 55.7% | 42.4% |
| Heroin | 2.9% | 38.4% | 33.3% | 4.3% | 1.6% |
| Cocaine | 7.8% | 57.6% | 60.6% | 17.8% | 10.2% |
| Marijuana | 22.5% | 56.6% | 56.6% | 28.6% | 17.5% |
| Alcohol | 48.0% | 62.2% | 60.4% | 43.2% | 30.5% |
| Amphetamines | 5.5% | 52.3% | 54.5% | 12.9% | 7.5% |
| Inhalants | 6.5% | 8.6% | 6.1% | 3.0% | 1.0% |
| Other drugs | 4.4% | 13.0% | 13.3% | 4.8% | 2.2% |

African American

| | M/MS | IDU | | F/MS | |
|--------------|-------|-------|--------|-------|--------|
| | Male | Male | Female | Male | Female |
| Drug use | 64.7% | 96.5% | 96.7% | 71.5% | 46.8% |
| Heroin | 1.2% | 46.5% | 48.9% | 7.1% | 3.8% |
| Cocaine | 18.8% | 80.2% | 79.3% | 30.3% | 19.9% |
| Marijuana | 34.1% | 57.0% | 48.9% | 44.6% | 24.1% |
| Alcohol | 50.6% | 71.5% | 64.1% | 55.5% | 31.0% |
| Amphetamines | 1.2% | 20.3% | 14.1% | 3.2% | 1.4% |
| Inhalants | 3.5% | 3.5% | 1.1% | 0.5% | 0.3% |
| Other drugs | 1.2% | 3.5% | 4.3% | 1.2% | 0.4% |

Hispanic

| | M/MS | IDU | | F/MS | |
|--------------|-------|-------|--------|-------|--------|
| | Male | Male | Female | Male | Female |
| Drug use | 62.0% | 86.1% | 85.7% | 67.3% | 40.4% |
| Heroin | 3.3% | 39.2% | 40.0% | 4.5% | 0.6% |
| Cocaine | 10.9% | 59.5% | 62.9% | 27.0% | 12.4% |
| Marijuana | 22.8% | 59.5% | 40.0% | 35.6% | 19.1% |
| Alcohol | 54.3% | 67.1% | 54.3% | 57.4% | 33.4% |
| Amphetamines | 5.4% | 39.2% | 28.6% | 8.2% | 2.9% |
| Inhalants | 8.7% | 15.2% | 5.7% | 3.6% | 2.5% |
| Other drugs | 5.4% | 8.9% | 8.6% | 2.9% | 2.5% |

Dallas**White**

| | M/MS | IDU | | F/MS | |
|--------------|-------------|-------------|---------------|-------------|---------------|
| | Male | Male | Female | Male | Female |
| Drug use | 59.0% | 92.5% | 91.4% | 59.5% | 54.2% |
| Heroin | 1.2% | 32.3% | 37.9% | 2.2% | 2.1% |
| Cocaine | 13.4% | 47.7% | 45.0% | 15.8% | 11.3% |
| Marijuana | 17.5% | 25.2% | 24.6% | 22.2% | 17.9% |
| Alcohol | 50.9% | 47.0% | 42.5% | 39.8% | 34.2% |
| Amphetamines | 7.4% | 24.1% | 23.9% | 8.4% | 4.6% |
| Inhalants | 14.1% | 5.3% | 2.1% | 2.4% | 0.7% |
| Other drugs | 8.8% | 13.5% | 9.3% | 4.9% | 3.6% |

African American

| | M/MS | IDU | | F/MS | |
|--------------|-------------|-------------|---------------|-------------|---------------|
| | Male | Male | Female | Male | Female |
| Drug use | 56.1% | 90.4% | 80.8% | 64.9% | 42.3% |
| Heroin | 2.3% | 31.5% | 30.8% | 0.9% | 0.5% |
| Cocaine | 9.2% | 38.4% | 36.7% | 24.3% | 10.3% |
| Marijuana | 11.0% | 13.7% | 20.8% | 23.3% | 13.7% |
| Alcohol | 40.5% | 23.3% | 30.8% | 36.1% | 26.4% |
| Amphetamines | 3.5% | 2.7% | 1.7% | 0.6% | 0.8% |
| Inhalants | 5.8% | | | 0.2% | 0.3% |
| Other drugs | 4.0% | 8.2% | 2.5% | 1.4% | 0.9% |

Hispanic

| | M/MS | IDU | | F/MS | |
|--------------|-------------|-------------|---------------|-------------|---------------|
| | Male | Male | Female | Male | Female |
| Drug use | 56.6% | 93.5% | 71.4% | 46.3% | 20.9% |
| Heroin | 1.9% | 45.2% | 39.3% | 2.3% | 0.8% |
| Cocaine | 16.9% | 41.9% | 28.6% | 11.3% | 4.1% |
| Marijuana | 19.1% | 22.6% | 17.9% | 16.7% | 6.2% |
| Alcohol | 50.6% | 58.1% | 32.1% | 33.5% | 16.9% |
| Amphetamines | 4.9% | 19.4% | 7.1% | 3.5% | 0.2% |
| Inhalants | 12.0% | 3.2% | | 2.7% | 0.3% |
| Other drugs | 6.0% | 9.7% | 3.6% | 1.2% | 0.3% |

North East Metroplex**White**

| | M/MS | IDU | | F/MS | |
|--------------|-------------|-------------|---------------|-------------|---------------|
| | Male | Male | Female | Male | Female |
| Drug use | 52.4% | 83.0% | 83.1% | 32.2% | 26.5% |
| Heroin | 1.0% | 27.0% | 22.0% | 1.2% | 1.4% |
| Cocaine | 8.6% | 37.0% | 32.2% | 8.3% | 4.8% |
| Marijuana | 14.3% | 39.0% | 32.2% | 13.6% | 11.6% |
| Alcohol | 45.7% | 46.0% | 45.8% | 24.5% | 18.6% |
| Amphetamines | 6.7% | 35.0% | 27.1% | 4.7% | 3.7% |
| Inhalants | 5.7% | 12.0% | 5.1% | 0.9% | 0.6% |
| Other drugs | 6.7% | 7.0% | 16.9% | 3.8% | 1.8% |

African American

| | M/MS | IDU | | F/MS | |
|--------------|-------------|-------------|---------------|-------------|---------------|
| | Male | Male | Female | Male | Female |
| Drug use | 50.0% | 77.8% | 60.0% | 39.5% | 26.6% |
| Heroin | | 22.2% | 40.0% | 2.6% | |
| Cocaine | 12.5% | 44.4% | 40.0% | 14.9% | 6.2% |
| Marijuana | 12.5% | 22.2% | 20.0% | 20.2% | 6.8% |
| Alcohol | 25.0% | 55.6% | 20.0% | 21.9% | 14.1% |
| Amphetamines | | 22.2% | 20.0% | 5.3% | |
| Inhalants | | | | 0.9% | |
| Other drugs | | | | 0.9% | 1.1% |

Hispanic

| | M/MS | IDU | | F/MS | |
|--------------|-------------|-------------|---------------|-------------|---------------|
| | Male | Male | Female | Male | Female |
| Drug use | 55.6% | 100.0% | 75.0% | 23.1% | 27.6% |
| Heroin | | | 25.0% | 1.5% | |
| Cocaine | 11.1% | 50.0% | 50.0% | 6.2% | 1.3% |
| Marijuana | 16.7% | 25.0% | 25.0% | 12.3% | 7.9% |
| Alcohol | 55.6% | 50.0% | 50.0% | 13.8% | 21.1% |
| Amphetamines | 5.6% | | 25.0% | 3.1% | 1.3% |
| Inhalants | 5.6% | | | 3.1% | |
| Other drugs | | | | 1.5% | 1.3% |

Rural North Texas**White**

| | M/MS | IDU | | F/MS | |
|--------------|-------------|-------------|---------------|-------------|---------------|
| | Male | Male | Female | Male | Female |
| Drug use | 66.0% | 82.9% | 92.0% | 56.0% | 43.2% |
| Heroin | | 15.9% | 13.9% | 2.2% | 0.8% |
| Cocaine | 8.3% | 39.5% | 46.3% | 11.0% | 5.3% |
| Marijuana | 20.8% | 37.4% | 38.3% | 21.8% | 12.1% |
| Alcohol | 59.0% | 42.8% | 45.8% | 40.3% | 31.9% |
| Amphetamines | 2.8% | 46.7% | 54.2% | 5.9% | 3.9% |
| Inhalants | | 1.8% | | 1.2% | 0.5% |
| Other drugs | 2.1% | 16.5% | 15.4% | 6.6% | 3.2% |

African American

| | M/MS | IDU | | F/MS | |
|--------------|-------------|-------------|---------------|-------------|---------------|
| | Male | Male | Female | Male | Female |
| Drug use | 80.0% | 81.8% | 100.0% | 58.8% | 33.6% |
| Heroin | | 18.2% | 20.0% | 0.6% | |
| Cocaine | 20.0% | 54.5% | 80.0% | 18.6% | 16.8% |
| Marijuana | 20.0% | 9.1% | 20.0% | 22.0% | 16.0% |
| Alcohol | 80.0% | 18.2% | 20.0% | 41.8% | 26.1% |
| Amphetamines | | 9.1% | 20.0% | 1.1% | 0.8% |
| Inhalants | | | | | |
| Other drugs | | 4.5% | 20.0% | 1.7% | 0.8% |

Hispanic

| | M/MS | IDU | | F/MS | |
|--------------|-------------|-------------|---------------|-------------|---------------|
| | Male | Male | Female | Male | Female |
| Drug use | 59.1% | 77.8% | 87.5% | 73.7% | 37.5% |
| Heroin | | 8.3% | 12.5% | 1.8% | |
| Cocaine | 13.6% | 44.4% | 12.5% | 25.4% | 9.7% |
| Marijuana | 22.7% | 38.9% | 12.5% | 40.4% | 8.3% |
| Alcohol | 45.5% | 47.2% | 25.0% | 57.9% | 19.4% |
| Amphetamines | | 33.3% | | 8.8% | 2.8% |
| Inhalants | 4.5% | | | 3.5% | 1.4% |
| Other drugs | | 8.3% | | 5.3% | 2.8% |